
TESTBARK

MATERNAL-NEWBORN NURSING: THE CRITICAL COMPONENTS OF NURSING CARE 3RD EDITION DURHAM CHAPMAN



TEST BARK

Maternal-Newborn Nursing: The Critical Components of Nursing Care 3rd Edition Durham Chapman Test Bank

Table of Content

19. Alterations in Women's Health

Maternity Nursing Overview
1. Trends and Issues
2. Ethics and Standards of Practice Issues
The Antepartal Period
3. Genetics, Conception, Fetal Development, and Reproductive Technology
4. Physiological Aspects of Antepartum Care
5. The Psycho-Social-Cultural Aspects of the Antepartum Period
6. Antepartal Tests
7. High-Risk Antepartum Nursing Care
Intrapartal Period
8. Intrapartum Assessment and Interventions
9. Fetal Heart Rate Assessment
10. High-Risk Labor and Birth
11. Intrapartum and Postpartum Care of the Cesarean Birth Families
Postpartal Period
12. Postpartum Physiological Assessments and Nursing Care
13. Transition to Parenthood
14. High-Risk Postpartum Nursing Care
Neonatal Period
15. Physiological and Behavioral Responses of the Neonate
16. Discharge Planning and Teaching
17. High-Risk Neonatal Nursing Care
Women's Health
18. Well Women's Health

MULTIPLE CHOICE

- 1. The nurse is caring for a patient who is in labor with her first child. The patient's mother is present for support and notes that things have changed in the delivery room since she last gave birth in the early 1980s. Which current trend or intervention may the patient's mother find most different?
 - 1. Fetal monitoring throughout labor
 - 2. Postpartum stay of 10 days
 - 3. Expectant partner and family in operating room for cesarean birth
 - 4. Hospital support for breastfeeding

ANS: 4

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 1. Discuss current trends in the management of labor and birth

Page: 4

Heading: Table 1-1: Past and Present Trends

Integrated Processes: Nursing Process

Client Need: Health Promotion and Maintenance

Cognitive Level: Application [Applying]

Concept: Evidence-Based Practice

Difficulty: Moderate

	Feedback
1	This is incorrect. Fetal monitoring during labor began in the late 1970s. As such,
	this likely would have occurred during the mother's labor and delivery during
	the 1980s.
2	This is incorrect. In the past, the average hospital postpartum stay was 10 days.
	Presently, the average postpartum stay is 48 hours or less.
3	This is incorrect. In the past, expectant partners and families were excluded from
	the labor and birth experience. Present trends involve the expectant partner and
	family in the labor and birth experience, including presence in the operating
	room for cesarean births.
4	This is correct. Hospital support for breastfeeding, including a lactation
	consultant and employment of the Baby-Friendly Hospital Initiative, were both
	enacted during the early 1990s.

PTS: 1 CON: Evidence-Based Practice

- 2. A patient with a history of hypertension is giving birth. During delivery, the staff was not able to stabilize the patient's blood pressure. As a result, the patient died shortly after delivery. This is an example of what type of death?
 - 1. Early maternal death
 - 2. Late maternal death
 - 3. Direct obstetric death
 - 4. Indirect obstetric death

ANS: 4

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 2. Discuss current trends in maternal and infant health

outcomes. Page: 7

Heading: Trends > Maternal Death and Mortality Rates

Integrated Processes: Nursing Process

Client Need: Physiological Integrity: Reduction of Risk Potential

Cognitive Level: Application [Applying]

Concept: Ante/Intra/Post-partum

Difficulty: Hard

	Feedback
1	This is incorrect. Early maternal death is not an example of maternal death.
	Examples of maternal death include late maternal death, indirect obstetric death,
	direct obstetric death, and pregnancy-related death.
2	This is incorrect. Late maternal death occurs 42 days after termination of
	pregnancy from a direct or indirect obstetric cause.
3	This is incorrect. Direct obstetric death results from complications during
	pregnancy, labor, birth, and/or postpartum period.
4	This is correct. Indirect obstetric death is caused by a preexisting disease, or a
	disease that develops during pregnancy.

PTS: 1 CON: Ante/Intra/Post-partum

- 3. The nurse is providing education to a patient who has given birth to her first child and is being discharged home. The patient expressed concern regarding infant mortality and sudden infant death syndrome (SIDS). The patient had an uncomplicated pregnancy, labor, and vaginal delivery. She has a body mass index of 25 and has no other health conditions. The infant is healthy and was delivered full-term. What will be most helpful thing to explain to the patient?
 - 1. Uses of extracorporeal membrane oxygenation therapy (ECMO)
 - 2. Uses of exogenous pulmonary surfactant
 - 3. The Baby-Friendly Hospital Initiative
 - 4. The Safe to Sleep campaign

ANS: 4

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 3. Identify leading causes of infant death.

Page: 7

Heading: Trends > Infant Mortality Rates Integrated Processes: Nursing process

Client Need: Safe and Effective Care Environment: Safety and Infection Control

Cognitive Level: Application [Applying]

Concept: Health Promotion

Difficulty: Moderate

	Feedback
1	This is incorrect. EMCO has been cited as one of the factors that has reduced
	infant mortality among preterm infants.

2	This is incorrect. Although advances in medical treatments have decreased infant
	mortality, exogenous pulmonary surfactant is primarily used to reduce mortality of
	preterm infants.
3	This is incorrect. The Baby-Friendly Hospital Initiative was developed to support
	breastfeeding and is not directly linked to reduced infant mortality or SIDS.
4	This is correct. The Back to Sleep campaign and the Safe to Sleep campaigns were
	designed to promote healthy infant sleeping habits. The decrease in SIDS from
	1995 to 2015 was attributed to the Safe to Sleep campaign.

PTS: 1 **CON: Health Promotion**

- 4. The nurse is caring for a 14-year-old patient who is 32 weeks pregnant. After complaining of genital sores and discomfort, the patient tests positive for syphilis. The fetus is at increased risk of which condition?
 - 1. Diabetes
 - 2. Blindness
 - 3. Pneumonia
 - 4. Hypertension

ANS: 4

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 4. Discuss current maternal and infant health issues.

Page: 10

Heading: Issues > Teen Pregnancy > Implications of Teen Pregnancy and Birth

Integrated Processes: Nursing Process

Client Need: Physiological Integrity: Reduction of Risk Potential

Cognitive Level: Application [Applying]

Concept: Ante/Intra/Post-partum

Difficulty: Moderate

	Feedback
1	This is incorrect. Maternal obesity increases a child's risk of developing
	childhood obesity and diabetes.
2	This is correct. Neonatal blindness, maternal death, and neonatal death are all
	associated with a patient who contracts syphilis during pregnancy.
3	This is incorrect. Chlamydial pneumonia is associated with maternal chlamydia.
4	This is incorrect. Teen mothers may have a higher risk of contracting sexually
	transmitted illnesses and hypertension during pregnancy; however, maternal
	syphilis is not associated with fetal hypertension.

PTS: 1 CON: Ante/Intra/Post-partum

- 5. The nurse is caring for a 15-year-old female who is pregnant with her first child. In her previous prenatal visit, the patient tested negative for chlamydia, syphilis, gonorrhea, and HIV. Based on the information provided, which condition is the patient's baby at higher risk for?
 - 1. Intestinal problems
 - 2. Neonatal conjunctivitis
 - 3. Blindness

4. Pneumonia

ANS: 1

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 4. Discuss current maternal and infant health issues.

Page: 11

Heading: Issues > Teen Pregnancy Integrated Processes: Nursing Process

Client Need: Physiological Integrity: Reduction of Risk Potential

Cognitive Level: Application [Applying]

Concept: Health Promotion

Difficulty: Difficult

	Feedback
1	This is correct. Infants born to teen mothers are at increased risk for various conditions related to prematurity, including infant death, intestinal problems, and/or respiratory distress syndrome.
2	This is incorrect. Infants born to teen mothers who have gonorrhea are at increased risk of neonatal conjunctivitis and blindness.
3	This is incorrect. Infants born to teen mothers with syphilis and gonorrhea are at increased risk of blindness.
4	This is incorrect. Infants born to teen mothers with chlamydia may be at increased risk of developing chlamydial pneumonia.

PTS: 1 CON: Health Promotion

- 6. The nurse is caring for a 23-year-old patient who arrives at the clinic for a pregnancy test. The test confirms the patient is pregnant. The patient states, "I do not need to stop smoking my electronic cigarette because it will not harm my baby." Which is the best response by the nurse?
 - 1. "You are correct. Electronic cigarettes are not harmful during pregnancy."
 - 2. "Tobacco products, including electronic cigarettes, should not be used during pregnancy due to risking nicotine toxicity."
 - 3. "According to the FDA, although electronic cigarettes are safe for you, they can cause harm to the fetus during pregnancy."
 - 4. "Electronic cigarettes are considered harmful only in the first trimester."

ANS: 2

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 4. Discuss current maternal and infant health issues.

Page: 12

Heading: Issues > Tobacco and Electronic Cigarette Use During Pregnancy

Integrated Processes: Nursing Process

Client Need: Safe and Effective Care Environment: Safety and Infection Control

Cognitive Level: Application [Applying]

Concept: Ante/Intra/Post-partum

Difficulty: Moderate

Feedback

1	This is incorrect. Electronic cigarettes can be harmful during pregnancy.
2	This is correct. Pregnant women should not use tobacco products or electronic
	cigarettes during pregnancy.
3	This is incorrect. Electronic cigarettes are not controlled by the FDA and may be
	harmful to both mother and fetus.
4	This is incorrect. Electronic cigarettes are considered harmful during pregnancy.

PTS: 1 CON: Ante/Intra/Post-partum

- 7. The nurse is caring for a 16-year-old patient who is 32 weeks pregnant with her first child, who is male. The patient's mother has accompanied her to today's visit. During the nursing assessment, the patient mentions that she is no longer in a relationship with the baby's father but her mother plans to help her. However, the patient's mother asks whether this will have any impact on the child. Which should the nurse indicate the child is at increased risk of during his adolescence?
 - 1. Hypertension
 - 2. Diabetes
 - 3. Alcohol abuse
 - 4. Intraventricular bleeding

ANS: 3

Chapter: Chapter 1 Trends and Issues

Chapter Learning Objective: 4. Discuss current maternal and infant health issues.

Page: 12

Heading: Issues > Teen Pregnancy > Implications of Teen Pregnancy and Birth

Integrated Processes: Nursing Process

Client Need: Physiological Integrity: Reduction of Risk Potential

Cognitive Level: Application [Applying]

Concept: Ante/Intra/Post-partum

Difficulty: Moderate

	Feedback
1	This is incorrect. Teen mothers, rather than their children, are at increased risked
	of hypertension during pregnancy.
2	This is incorrect. Children born to mothers who are obese have an increased risk
	of developing childhood obesity and childhood diabetes.
3	This is correct. Statistics have shown that adolescent boys without an involved
	father may be at higher risk of incarceration, dropping out of school, and abusing
	drugs or alcohol.
4	This is incorrect. Children born to teen mothers are at increased risk for health
	problems associated with low birth weight, including intraventricular bleeding.

PTS: 1 CON: Ante/Intra/Post-partum

- 8. The nurse is caring for a patient at 7 weeks gestation. The nurse suspects that a pregnant patient may have been using marijuana. With consent, the nurse confirms via urine drug screen. Which statement by the nurse is most appropriate?
 - 1. "Did you smoke marijuana when pregnant with your other child?"
 - 2. "To avoid negative effects on your baby, you'll need to stop using marijuana